



# Sclerochronological study of the gigantic inoceramids *Sphenocerasmus schmidtii* and *S. sachalinensis* from Hokkaido, northern Japan

ERIC O. WALLISER , KAZUSHIGE TANABE, YOSHINORI HIKIDA, KOTARO SHIRAI AND BERND R. SCHÖNE

## LETHAIA



Walliser E. O., Tanabe K., Hikida Y., Shirai K., & Schöne B. R. 2019: Sclerochronological study of the gigantic inoceramids *Sphenocerasmus schmidtii* and *S. sachalinensis* from Hokkaido, northern Japan. *Lethaia*, Vol. 52, pp. 410–428.

Here, we present the first sclerochronological investigation of shells of the gigantic inoceramids *Sphenocerasmus schmidtii* and *S. sachalinensis* from the middle Campanian cold seep carbonate-bearing strata of the Yezo Basin in Hokkaido (northern Japan). Stable carbon ( $\delta^{13}\text{C}$ ) and oxygen ( $\delta^{18}\text{O}$ ) isotope values were measured in the aragonitic and calcitic shell layers of both species and compared to those of other co-occurring benthic (mainly bivalves and gastropods) and demersal molluscs (ammonites). Sedimentological and stable isotope data suggest that these bivalves lived near cold seeps and were exposed to high  $\text{H}_2\text{S}$  level in the seawater. The inoceramid shells exhibited higher  $\delta^{13}\text{C}$  and lower  $\delta^{18}\text{O}$  values than the coeval non-cold seep molluscs. We ascribed the anomalous isotopic pattern to a combination of vital and environmental effects determined by the hosting of chemosymbionts and the exposure to warm interstitial waters. Inoceramid  $\delta^{13}\text{C}$  minima coincided with growth lines and likely reflect changes in nutrient supply by the chemosymbionts. Absolute temperatures estimated from  $\delta^{18}\text{O}$  values of *Sphenocerasmus schmidtii* and *S. sachalinensis* were, on average, ca. 4–5°C warmer than those reconstructed for the non-seepage environment ( $19.3 \pm 0.7^\circ\text{C}$ ). Short-term  $\delta^{18}\text{O}$  fluctuations of the inoceramid material indicate local temperature ranges of up to 5.2°C, that is four times larger than those reconstructed from the benthic and demersal fauna (1.3°C). In general, our data suggest that the stable carbon and oxygen isotope values of the studied *Sphenocerasmus* spp. were strongly affected by short-term fluctuations in seepage activity and do not reflect seasonal fluctuations. □ *Campanian, chemosymbiosis, cold seep, oxygen and carbon isotopes, sclerochronology.*

Eric O. Walliser  [walliser@uni-mainz.de], and Bernd R. Schöne [schoeneb@uni-mainz.de], Institute of Geosciences, University of Mainz, Johann-Joachim-Becher-Weg 21, 55128 Mainz, Germany; Kazushige Tanabe [tanabe@um.u-tokyo.ac.jp], The University Museum, The University of Tokyo, Hongo 7-3-1, Tokyo 113-0033, Japan; Yoshinori Hikida [nmhikida@coral.ocn.ne.jp], Nakagawa Museum of Natural History, Nakagawa (Hokkaido), 28-9 Yasukawa Nakagawa, Hokkaido Prefecture 098-2625, Japan; Kotaro Shirai [kshirai@aori.u-tokyo.ac.jp], Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Chiba Prefecture 277-0882, Japan; manuscript received on 16/07/2018; manuscript accepted on 7/12/2018.

Inoceramid shells provide an exceptional geochemical archive for Late Cretaceous high-resolution palaeoclimate reconstructions (Jiménez Berrocoso *et al.* 2008; Walliser *et al.* 2018). However, their  $\delta^{13}\text{C}$  values are usually higher, and sometimes (but not always),  $\delta^{18}\text{O}$  values are lower than expected from isotopic equilibrium fractionation with the Cretaceous seawater (Tourtelot & Rye 1969; Pirrie & Marshall 1990; MacLeod & Hoppe 1992; Ludvigson *et al.* 1994; Elorza & García-Garmilla 1996; Fisher & Arthur 2002; He *et al.* 2005; Henderson & Price 2012; Zakharov *et al.* 2012). Remarkably, inoceramids exhibit such stable isotope pattern when found embedded in  $^{13}\text{C}$ -depleted carbonate

concretions with high amounts of microcrystalline pyrite (Tourtelot & Rye 1969; Whittaker *et al.* 1987; Wright 1987; Carpenter *et al.* 1988; Pirrie & Marshall 1990; Ludvigson *et al.* 1994; Henderson & Price 2012; Zakharov *et al.* 2012). Such concretions are diagnostic features for bacterially mediated anoxic oxidation of methane (AOM) and are often associated with ancient cold seeps (Aharon & Fu 2000; Peckmann & Thiel 2004; Angeletti *et al.* 2015; Reitner *et al.* 2015), that is sites at which hydrocarbon-loaded fluids vent out of the seafloor.

Methane seepages are habitats that are depleted in  $^{12}\text{C}$  and enriched in  $\text{H}_2\text{S}$  which can only be colonized by a highly specialized chemosymbiotic fauna (Levin